

Laser and Fourier Transform Spectroscopy of Novel Propellant Molecules

Peter F. Bernath

Department of Chemistry
University of Waterloo
Waterloo, ON
Canada N2L 3G1

January 1994

Final Report


DTIC
ELECTE
MAR 18 1994
S B D

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

94-08774



20pg



PHILLIPS LABORATORY--
Propulsion Directorate
AIR FORCE MATERIEL COMMAND
EDWARDS AIR FORCE BASE CA 93524-7001

DTIC QUALITY ESTIMATED 1

94 3 18 0 42

NOTICE

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise, or in any way licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may be related thereto.

FOREWORD

This report was prepared by Department of Chemistry, University of Waterloo, Ontario, Canada, under contract F29601-91-C-0100, for Operating Location AC, Phillips Laboratory, Edwards AFB, CA. 93524-7001. Project Manager for Phillips Laboratory was Bob Behdadnia.

This report has been reviewed and is approved for release and distribution in accordance with the distribution statement on the cover and on the SF Form 298.



ROBERT BEHDADNIA
Project Manager



STEPHEN L. RODCERS
Chief, Emerging Technologies Branch



LEONARD C. BROLINE, Lt Col, USAF
Director,
Fundamental Technologies Division



RANNEY G. ADAMS 93-198
Public Affairs Director

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Attn and/or Special
A-1	

REPORT DOCUMENTATION PAGE			Form Approved OMB No 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions searching existing data sources gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0740-0188), Washington DC 20503.				
1. AGENCY USE ONLY (LEAVE BLANK)		2. REPORT DATE January 1994		3. REPORT TYPE AND DTAE COVERED Final 1 Apr 90 - 30 Sep 93
4. TITLE AND SUBTITLE Laser and Fourier Transform Spectroscopy of Novel Propellant Molecules			5. FUNDING NUMBERS C: F29601-91-C-0100 PE: 62302F PR: 3058 TA: 00DS	
6. AUTHOR(S) Peter F. Bernath				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry University of Waterloo Waterloo, Ontario Canada N2L 3G1			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Phillips Laboratory OLAC PL/RKFE 9 Antares Road Edwards AFB CA 93524-7680			10. SPONSORING/MONITORING AGENCY REPORT NUMBER PL-TR-93-3088	
11. SUPPLEMENTARY NOTES COSATI CODE(S):				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for Public Release; Distribution is unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (MAXIMUM 200 WORDS) A wide variety of molecules were studied by the techniques of Fourier transform spectroscopy and visible laser spectroscopy. Molecules studied include CCN, CaBH ₄ , CH ₃ N, CN, IF, ICl, IBr, BC, CH, CH, C ₂ , C ₃ , C ₆₀ , LiF, AlF, AlCl, HF, HCl, B ₃ F ₂ , BiH, AlH, BaH, and GaH. These molecules were detected by infrared, visible and ultraviolet emission and absorption measurements.				
14. SUBJECT TERMS visible; Fourier transform; spectroscopy; ultraviolet; laser; UV/VIS; interhalogen(s); metal(s); borohydrides; C ₂ ; C ₃ ; C ₆₀ ; dihalide(s); methyl nitrene; dicarbon nitride; CH; CCN			15. NUMBER OF PAGES 20	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT SAR	

TABLE OF CONTENTS

	Page
A. Introduction	1
B. Dicarbon Nitride, CCN	1
C. Metal Borohydrides, CaBH_4 , SrBH_4 , LiBH_4 and NaBH_4	1
D. Methyl Nitrene, CH_3N	2
E. Cyanogen, CN	2
F. Review Articles	2
G. Interhalogens, IF, ICl, IBr	5
H. Boron-Containing Diatomics, BC, BD, BH	5
I. CH	5
J. Pure Carbon Molecules, C_2 , C_3 and C_{60}	6
K. Metal Halides, LiF, AlF and AlCl	7
L. Hydrogen Halides, HF and HCl	7
M. Metal Dihalides, BeF_2	9
N. Metal Hydrides and Deuterides, BiH, BiD, AlH, AlD, BaH, GaH and GaD	9
O. Conclusions and Future Directions	10
P. Publications (supported by F29601-91-C-0100)	14

The goal of our work is the study of the spectroscopy and chemistry of a variety of energetic molecules. They have been studied by high resolution infrared and visible Fourier transform spectroscopy as well as visible laser spectroscopy. Many of these molecules such as C_2 , BC, BH and AlH may prove to be useful in boosting the specific impulse of the H_2/O_2 propellant system if they are trapped in solid hydrogen. Some of the molecules studied such as CH, CN, AlF, HCl, HF and AlCl are of interest in the modelling of current conventional aluminized propellants. Some of our work also has applications in the simulation of rocket plume signatures and in the Air Force chemical laser program.

B. Dicarbon Nitride, CCN.

The ultracold emission spectrum of CCN was observed with an Engelking-type [Engelking, *Rev. Sci. Instrum.* **57**, 2274 (1986)] free radical jet expansion source. The precursor diazoacetonitrile $HC(N_2)CN$ was seeded in He and discharged to provide CCN. The observation¹ of the 000-000, 000-001, 000-002 and 000-100 vibronic bands of the $\tilde{A}^2\Delta - \tilde{X}^2\Pi$ transition allowed the accurate determination of the ν_1 (1923.255 cm^{-1}), ν_3 (1050.764 cm^{-1}) and $2\nu_2$ (2094.816 cm^{-1}) vibrational frequencies. The very precise wavenumber scale of the Fourier transform spectrometer allows these infrared active vibrational modes to be determined from an electronic emission spectrum. Electronic emission spectra of ultracold free radicals detected by Fourier transform spectroscopy is a powerful technique for the studying of transient molecules.

C. Metal Borohydrides, $CaBH_4$, $SrBH_4$, $LiBH_4$ and $NaBH_4$.

Ca and Sr vapors react spontaneously with diborane (B_2H_6) to give $CaBH_4$ and $SrBH_4$ free radicals². We have detected the $\tilde{B}^2E - \tilde{X}^2A_1$ and $\tilde{A}^2A_1 - \tilde{X}^2A_1$ electronic transitions by low resolution laser-induced fluorescence. Some preliminary infrared emission experiments on $LiBH_4$

and NaBH_4 suggest that they can be vaporized in a furnace. The infrared spectra were very complex and there was considerable decomposition of the heated LiBH_4 and NaBH_4 solids.

D. Methyl Nitrene, CH_3N .

The near ultraviolet emission spectrum of methyl nitrene, CH_3N , was observed at a resolution of 0.07 cm^{-1} with a Fourier transform spectrometer. The $\bar{\text{A}}^3\text{E} - \bar{\text{X}}^3\text{A}_2$ system of CH_3N (analogous to the $\text{A}^3\Pi - \text{X}^3\Sigma^-$ system of NH) was rotationally analysed. The CH_3N molecule was made in an Engelking jet source using CH_3N_3 as the precursor. Our rotational constants were consistent with a short C-N bond length of 1.420 \AA . This short bond length was not predicted by the early ab initio calculations but is in agreement with the most recent results of H.F. Schaefer's group [Y. Xie et al. *J. Am. Chem. Soc.* 111, 5181 (1989)].


E. Cyanogen, CN .

In the same experiments which provided the CCN and CH_3N spectra, new spectra of CN were observed. Both the $\text{B}^2\Sigma^+ - \text{X}^2\Sigma^+$ transition⁴ (violet system) and the $\text{A}^2\Pi - \text{X}^2\Sigma^+$ transition⁵ (red system) were observed. These spectra were rotationally cold but vibrationally hot. The great simplification provided by the jet source compared to a conventional discharge is illustrated in Figure 1. The analysis of numerous vibrational bands allowed new potential energy curves to be calculated. These curves (Figure 2) extend to high v and are very useful for the calculation of the spectral and thermochemical properties of CN in high temperature environments. The CN molecule is an important free radical in the combustion of conventional propellants.

F. Review Articles.

Two review articles were published on the research supported by our Air Force contract. A review of the high resolution infrared spectra of transient molecules was published in the

Figure 1: The 0-0 band of the $B^2\Sigma^+ - X^2\Sigma^+$ transition of CN observed at room temperature (Cossart, top) and 90K (jet, bottom).






Figure 2: The RKR potential energy curves of the $X^2\Sigma^+$, $A^2\Pi$ and $B^2\Sigma^+$ electronic states of CN.

Annual Review of Physical Chemistry⁶. While a review of our organometallic research on molecules such as CaBH_4 was published in Science magazine⁷.

G. Interhalogens, IF, ICl and IBr.

The vibration-rotation spectra of IF,⁸ ICl⁹ and IBr¹⁰ were recorded. Interhalogens are not of direct interest to the HEDM program (except as oxidizers), but they are important to the Air Force chemical laser program. We recorded the ICl and IBr spectra to test the far infrared performance of our new Bruker Fourier transform spectrometer. Surprisingly no high resolution vibration-rotation spectra of ICl and IBr were available in the literature prior to our work. The band origins at 384 cm^{-1} for ICl and 267 cm^{-1} for IBr are in the far infrared where conventional spectrometers have difficulty operating.

H. Boron-Containing Molecules, BC, BH and BD.

The BC molecule was discovered¹¹ by detection of the $\text{B}^4\Sigma^- - \text{X}^4\Sigma^-$ transition in a composite wall ($\text{B}_4\text{C}/\text{Cu}$) hollow cathode sputter source. The electronic emission of BC near 5590 \AA was recorded with the Kitt Peak Fourier transform spectrometer. As a bonus, the same spectra contained the $\text{A}^1\Pi - \text{X}^1\Sigma^+$ transition of BH. By adding D_2 to the source, the corresponding transition of BD was observed for both ^{10}B and ^{11}B isotopes.¹² The BC and BH molecules are both important high energy density materials (HEDM) since they yield (P. Carrick, private communication) a substantial improvement in the specific impulse (I_{sp}) of the hydrogen/oxygen system if they are trapped in solid hydrogen.

I. CH.

New emission measurements of the $\text{A}^2\Delta - \text{X}^2\Pi$ and $\text{B}^2\Sigma^- - \text{X}^2\Pi$ electronic transitions as well as the infrared vibration-rotation transitions were recorded as impurities in spectra recorded

for other purposes. A simultaneous analysis of all available data¹³ has resulted in a greatly improved set of molecular constants for CH. The CH molecule, like BH, is potential additive to solid hydrogen. In addition CH is a key free radical when any hydrocarbons (i.e. conventional fuels) are burned. Our new constants in the ground state cover the high-J and high-v levels necessary for simulations of the hot CH molecule during the combustion of propellants.

J. Pure Carbon Molecules, C_2 , C_3 and C_{60} .

Pure carbon molecules continue to fascinate the scientific community. Although C_{60} is too stable to be useful a high energy density material, C_2 , C_3 and smaller clusters are more relevant. For instance, C_2 and C_3 can be synthesized by trapping carbon atoms in solid hydrogen. The accidental or intentional agglomeration of the C atoms leads to C_2 and C_3 .

A new metastable state of C_3 ($\bar{a}^3\Pi_u$) was discovered by observing the $\bar{b}^3\Pi_g - \bar{a}^3\Pi_u$ electronic transition in the infrared.¹⁴ The infrared transition of C_3 was detected near 6500 cm^{-1} in emission from a microwave discharge of hydrocarbons. The same transition was also seen in absorption with a diode laser since the lower $\bar{a}^3\Pi_u$ state has a long lifetime (20 ms in an argon matrix).

For C_{60} the gas-phase vibration-rotation bands were also seen in emission by vaporizing a sample of C_{60} . The hot C_{60} vapor showed¹⁵ the four infrared active bands expected on the basis of the previous solid state infrared work. This work demonstrates that our favourite technique of infrared emission spectroscopy also works well for large molecules in which the rotational structure is not resolved.

In the same Fourier transform spectra that yielded data for CH_3N , CCN and CN molecules, the C_2 Swan system, $d^3\Pi_g - a^3\Pi_u$, was also identified. As for CN , the use of the

Engelking jet source gave vibrationally hot but rotationally cold (90 K) spectra. These data were combined with some additional laser data taken from the literature to provide greatly improved constants for the Swan system of C_2 . Like CH and CN, C_2 is always present during the combustion of conventional propellants and is a potential HEDM molecule.

K. Metal Halides, LiF, AlF and AlCl.

We have found high resolution vibration-rotation emission spectroscopy to be a powerful tool for the spectroscopic characterization of high temperature materials. This technique has received little attention at long wavelengths (greater than 5 microns). If lithium or aluminum compounds are used as propellants with an oxidizer containing fluorine or chlorine then LiF, AlF and AlCl will be among the exhaust products. We have recorded the vibration-rotation emission spectra of LiF¹⁶ and AlF¹⁷ near 900 cm^{-1} as well as AlCl¹⁸ near 500 cm^{-1} . The quality of the spectra is illustrated (Figure 3) with a section of the AlCl spectrum near the 2-1 band head. Our data on AlCl has already been used to simulate the AlCl features detected in a rocket plume in the ultraviolet (J. Drakes, Arnold AFB, private communication).

L. Hydrogen Halides, HF and HCl.

During the course of our work with light metal halides, we observed the pure rotational emission spectra of hot HF^{19,20} and hot HCl²⁰. Our emission technique is thus useful for infrared electronic spectra (e.g. C_3) infrared vibration-rotation spectra of small molecules (e.g. AlH) as well as large molecules (e.g. C_{60}) and for the pure rotation spectra of small molecules (e.g. HCl). Indeed we find it to be the technique of choice for high temperature systems in general.

The vibration-rotation emission spectra of hot HCl and hot HF were recorded by the pyrolysis of the freon CHF_2Cl at 1000°C . Surprisingly the high-J and high-v transitions of HCl




Figure 3: The vibration-rotation emission spectrum of AlCl .

and HF were poorly known before our work. We measured these lines to a precision of better than $\pm 0.0002 \text{ cm}^{-1}$. These data are very useful for the simulation of the spectra of exhaust plumes of any propellant containing chlorine or fluorine.

M. Beryllium Difluoride, BeF_2 .

The technique of infrared emission spectroscopy can also be applied to polyatomic molecules such as BeF_2 . If Be is used as an additive to boost the performance of a propellant for space-based applications, then BeF_2 will be an exhaust product with fluorine oxidizers. We recorded the vibration-rotation emission spectrum²¹ of the antisymmetric stretching mode near 1550 cm^{-1} . The fundamental and numerous hot bands were analysed and an equilibrium molecular structure was determined [$r_e(\text{Be-F}) = 1.372971 \text{ \AA}$]. This work is the first complete analysis of a metal dihalide spectrum.

N. Metal Hydrides and Deuterides, BiH, BiD, AlH, AlD, BaH, GaH and GaD.

Our infrared emission work was extended to metal hydrides by observing BiH²², BiD²² and BaH²³. For technical reasons (ease of handling and favorable chemistry) we chose to start with Bi and Ba metals. When these experiments were successful we began a series of experiments on the Al + H₂ system.

Molten aluminum is very corrosive and we had some initial difficulties with the destruction of our furnace tubes. The solution to our materials problem was to use a carbon liner for our tube furnace. In this way we could operate up to 1600°C without difficulty. Excellent spectra of AlH (Figure 4 and 5) were recorded²⁴. No evidence for the formation of AlH₂ could be found in our spectra.

We have also begun to develop new data reduction techniques. In order to derive the best Al-H interaction potential we fitted our data directly to the eigenvalues of the appropriate Schrödinger equation with a parameterized potential ("inverse perturbation approach"). For our potential we used an extension of the modified Morse function suggested by Coxon [J. Coxon et al., Can J. Phys. 70, 40 (1992)]. The results of our fit are illustrated in Figure 6. This potential reproduces the experimental data available for the lower half of the well and is qualitatively correct for the upper half where only ab initio data (points in Figure 6) are available. These potentials are necessary to simulate the interaction of aluminum with hydrogen in an advanced propellant. In order to test our theoretical approach some spectra²⁵ of GaH and GaD were also recorded.

O. Conclusion and Future Directions

The techniques of laser and Fourier transform spectroscopy are able to characterize energetic molecules suitable as propellants. Infrared emission spectroscopy of high temperature molecules has proved to be a particularly useful technique. The molecules of interest can be made by a variety of techniques including electrical discharges and high temperature vaporization and reaction.

The performance of a potential propellant depends both on the reaction exothermicity and the molecular weight of the product molecules. Light elements are most suitable so that we have concentrated our work on elements such as Li, Be, B and Al. We believe that simple thermal evaporation in a high temperature furnace followed by a chemical reaction and deposition into a solid hydrogen matrix holds considerable promise as a scheme to increase the performance of the hydrogen/oxygen system. We recommend the study of the reactions of laser and thermally

Fig. 4 - Overview Emission Spectrum Of AlH




Fig. 5 - Band Head Of The Fundamental Band Of AlH





Figure 6: The AlH potential energy curve. The points are from a recent ab initio calculation.

Born-Oppenheimer Potential AlH



generated B, C and Al atoms with each other as well as with small molecules such as H₂.

The first step in the development of an advanced propellant system is the identification of suitable new atomic and molecular fuels and oxidizers. Ultimately these light, energetic molecules will have to be synthesized efficiently and stored in condensed phases to make a viable propellant. Our work reported here concentrates on the initial steps in the process of advanced propellant design.

P. Publications (supported by grant F29601-91-C-0100).

1. N. Oliphant, A. Lee, P.F. Bernath and C.R. Brazier, Fourier Transform Emission Spectroscopy of the Jet-Cooled CCN Free Radical, *J. Chem. Phys.* **92**, 2244-2247 (1990).
2. F.S. Pianalto, A.M.R.P. Bopegedera, W.T.M.L. Fernando, R. Hailey, L.C. O'Brien, C.R. Brazier, P.C. Keller and P.F. Bernath, Gas Phase Inorganic Chemistry: Laser Spectroscopy of Calcium and Strontium Monoborohydrides, *J. Am. Chem. Soc.* **112**, 7900-7903 (1990).
3. C.R. Brazier, P.G. Carrick and P.F. Bernath, Rotational Analysis of the 0^0_0 Band of the $\bar{A}^3E-\bar{X}^3A_2$ System of Methylnitrene, *J. Chem. Phys.* **96**, 919-926 (1992).
4. C.V.V. Prasad, P.F. Bernath, C. Frum and R. Engleman, Fourier Transform Jet Spectroscopy of the $B^2\Sigma^+ - X^2\Sigma^+$ Transition of CN, *J. Mol. Spectrosc.* **151**, 459-473 (1992).
5. C.V.V. Prasad and P.F. Bernath, Fourier Transform Jet Emission Spectroscopy of the $A^2\Pi-X^2\Sigma^+$ Transition of CN, *J. Mol. Spectrosc.* **156**, 327-340 (1992).
6. P.F. Bernath, High-Resolution Infrared Spectroscopy of Transient Molecules, *Ann. Rev. Phys. Chem.* **41**, 91-122 (1990).

7. P. Bernath, Gas-Phase Inorganic Chemistry: Monovalent Derivatives of Calcium and Strontium, *Science* **254**, 665-670 (1991).
8. C.I. Frum, R. Engleman, Jr. and P.F. Bernath, Fourier Transform Detection of the Vibration-Rotation Bands of IF, *Chem. Phys. Lett.* **167**, 356-361 (1990).
9. H.G. Hedderich, P.F. Bernath and G.A. McRae, The High Resolution Infrared Spectrum of Iodine Monochloride, *J. Mol. Spectrosc.* **155**, 384-392 (1992).
10. J.M. Campbell and P.F. Bernath, Vibration-Rotation Spectrum of Iodine Monobromide, *J. Mol. Spectrosc.* **158**, 339-346 (1993).
11. W.T.M.L. Fernando, L.C. O'Brien and P.F. Bernath, Fourier Transform Emission Spectroscopy: The $B^4\Sigma^- - X^4\Sigma^-$ Transition of BC, *J. Chem. Phys.*, **93**, 8482-8487 (1990).
12. W.T.M.L. Fernando and P.F. Bernath, Fourier Transform Spectroscopy of the $A^1\Pi - X^1\Sigma^+$ Transition of BH and BD, *J. Mol. Spectrosc.*, **145**, 392-402 (1991).
13. P.F. Bernath, C.R. Brazier, T. Olsen, R. Hailey, W.T.M.L. Fernando, C. Woods and J.L. Hardwick, Spectroscopy of the CH Free Radical, *J. Mol. Spectrosc.* **147**, 16-26 (1991).
14. H. Sasada, T. Amano, C. Jarman and P.F. Bernath, A New Triplet Band System of C_3 : The $\tilde{b}^3\Pi_g - \tilde{a}^3\Pi_u$ Transition, *J. Chem. Phys.*, **94**, 2401-2407 (1991).
15. C.I. Frum, R. Engleman, Jr., H.G. Hedderich, P.F. Bernath, L.D. Lamb and D.R. Huffman, The Infrared Emission Spectrum of Gas Phase C_{60} (Buckminster fullerene), *Chem. Phys. Lett.*, **176**, 504-508 (1991).
16. H.G. Hedderich, C.I. Frum, R. Engleman, Jr., and P.F. Bernath, The Infrared Emission Spectra of LiF and HF, *Can. J. Chem.* **69**, 1659-1671 (1991).

17. H.G. Hedderich and P.F. Bernath, The Infrared Emission Spectrum of Gaseous AlF. *J. Mol. Spectrosc.* **153**, 73-80 (1992).
18. H.G. Hedderich, M. Dulick and P.F. Bernath, High Resolution Emission Spectroscopy of AlCl at 20 μ m, *J. Chem. Phys.* (in press).
19. H.G. Hedderich, K. Walker and P.F. Bernath, An Improved Set of Rotational Constants for HF, *J. Mol. Spectrosc.* **149**, 314-316 (1991).
20. R.B. LeBlanc, J.B. White and P.F. Bernath, High Resolution Emission Spectra of HCl and Hf, *J. Mol. Spectrosc.* (in press).
21. C.I. Frum, R. Engleman, Jr., and P.F. Bernath, Fourier Transform Emission Spectroscopy of BeF₂ at 6.5 m, *J. Chem. Phys.* **95**, 1435-1440 (1991).
22. H.G. Hedderich and P.F. Bernath, Infrared Emission Spectroscopy of BiH and BiD, *J. Mol. Spectrosc.* **158**, 170-176 (1993).
23. K.A. Walker, H.G. Hedderich and P.F. Bernath, High Resolution Vibration-Rotation Emission Spectroscopy of BaH, *Mol. Phys.* **78**, 577-589 (1993).
24. J.B. White, M. Dulick and P.F. Bernath, High Resolution Infrared Emission Spectra of AlH and AlD, *J. Chem. Phys.* (in press).
25. J.M. Campbell, M. Dulick, D. Klapstein, J.B. White and P.F. Bernath, Infrared Emission Spectra of GaH and GaD, *J. Chem. Phys.* (in press).